

Konstanter Benommenheitsschwindel in aufrechter Körperhaltung

Persistent Postural-Perceptual Dizziness (PPPD)

Zeitschrift für Physiotherapeuten, April 2021, Stefan Schädler

Literaturliste

1. Bisdorff A, Von Brevern M, Lempert T, Newman-Toker DE. Classification of vestibular symptoms: towards an international classification of vestibular disorders. *J Vestib Res.* 2009;19(1-2):1-13.
2. Staab JP, Eckhardt-Henn A, Horii A, Jacob R, Strupp M, Brandt T, et al. Diagnostic criteria for persistent postural-perceptual dizziness (PPPD): Consensus document of the committee for the Classification of Vestibular Disorders of the Bárány Society. *J Vestib Res.* 2017;27(4):191-208.
3. Brandt T, Dieterich M. Phobischer Attacken Schwankschwindel, ein neues Syndrom? *Munch Med Wschr.* 1986;28:247-50.
4. Bronstein AM. Visual vertigo syndrome: clinical and posturography findings. *J Neurol Neurosurg Psychiatry.* 1995;59(5):472-6.
5. Bronstein AM. The visual vertigo syndrome. *Acta Otolaryngol Suppl.* 1995;520 Pt 1:45-8.
6. Staab JP, Ruckenstein MJ, Amsterdam JD. A prospective trial of sertraline for chronic subjective dizziness. *Laryngoscope.* 2004;114(9):1637-41.
7. Jacob RG, Woody SR, Clark SB, Lilienfeld SO, Hirsch BE, Kucera GD, et al. Discomfort with space and motion: a possible marker of vestibular dysfunction assessed by the situational characteristics questionnaire. *Journal of Psychopathology and Behavioral Assessment.* 1993;15(4):299-324.
8. Popkirov S, Staab JP, Stone J. Persistent postural-perceptual dizziness (PPPD): a common, characteristic and treatable cause of chronic dizziness. *Pract Neurol.* 2018;18(1):5-13.
9. DGN/ÖGN. Leitlinien Schwindel - Diagnostik. Österreichische Gesellschaft für Neurologie, Deutsche Gesellschaft für Neurologie. 2015;Kap. 051:1-26.
10. Staab JP, Ruckenstein MJ. Expanding the differential diagnosis of chronic dizziness. *Arch Otolaryngol Head Neck Surg.* 2007;133(2):170-6.
11. Staab JP. Chronic subjective dizziness. *Continuum (Minneap Minn).* 2012;18(5 Neuro-otology):1118-41.
12. Balaban CD, Jacob RG, Furman JM. Neurologic bases for comorbidity of balance disorders, anxiety disorders and migraine: neurotherapeutic implications. *Expert Rev Neurother.* 2011;11(3):379-94.
13. Cousins S, Cutfield NJ, Kaski D, Palla A, Seemungal BM, Golding JF, et al. Visual dependency and dizziness after vestibular neuritis. *PLoS One.* 2014;9(9):e105426.
14. Staab JP. Persistent Postural-Perceptual Dizziness. *Semin Neurol.* 2020;40(1):130-7.
15. Sohsten E, Bittar RS, Staab JP. Posturographic profile of patients with persistent postural-perceptual dizziness on the sensory organization test. *J Vestib Res.* 2016;26(3):319-26.
16. Krawczyk S, Tietze S, Swoboda W, Valkovic P, Brandt T. Artificial neural network: a new diagnostic posturographic tool for disorders of stance. *Clin Neurophysiol.* 2006;117(8):1692-8.
17. Brandt T, Strupp M, Novozhilov S, Krawczyk S. Artificial neural network posturography detects the transition of vestibular neuritis to phobic postural vertigo. *J Neurol.* 2012;259(1):182-4.
18. Holmberg J, Tjernstrom F, Karlberg M, Fransson PA, Magnusson M. Reduced postural differences between phobic postural vertigo patients and healthy subjects during a postural threat. *J Neurol.* 2009;256(8):1258-62.
19. Querner V, Krawczyk S, Dieterich M, Brandt T. Patients with somatoform phobic postural vertigo: the more difficult the balance task, the better the balance performance. *Neurosci Lett.* 2000;285(1):21-4.
20. Querner V, Krawczyk S, Dieterich M, Brandt T. Phobic postural vertigo. Body sway during visually induced roll vection. *Exp Brain Res.* 2002;143(3):269-75.
21. Wuehr M, Brandt T, Schniepp R. Distracting attention in phobic postural vertigo normalizes leg muscle activity and balance. *Neurology.* 2017;88(3):284-8.

22. Indovina I, Riccelli R, Chiarella G, Petrolo C, Augimeri A, Giofre L, et al. Role of the Insula and Vestibular System in Patients with Chronic Subjective Dizziness: An fMRI Study Using Sound-Evoked Vestibular Stimulation. *Front Behav Neurosci.* 2015;9:334.
23. Riccelli R, Passamonti L, Toschi N, Nigro S, Chiarella G, Petrolo C, et al. Altered Insular and Occipital Responses to Simulated Vertical Self-Motion in Patients with Persistent Postural-Perceptual Dizziness. *Front Neurol.* 2017;8:529.
24. Huber J, Flanagan VL, Popp P, Zu Eulenburg P, Dieterich M. Network changes in patients with phobic postural vertigo. *Brain Behav.* 2020;10(6):e01622.
25. Lee JO, Lee ES, Kim JS, Lee YB, Jeong Y, Choi BS, et al. Altered brain function in persistent postural perceptual dizziness: A study on resting state functional connectivity. *Hum Brain Mapp.* 2018;39(8):3340-53.
26. Nigro S, Indovina I, Riccelli R, Chiarella G, Petrolo C, Lacquaniti F, et al. Reduced cortical folding in multi-modal vestibular regions in persistent postural perceptual dizziness. *Brain Imaging Behav.* 2019;13(3):798-809.
27. Wurthmann S, Naegel S, Schulte Steinberg B, Theysohn N, Diener HC, Kleinschnitz C, et al. Cerebral gray matter changes in persistent postural perceptual dizziness. *J Psychosom Res.* 2017;103:95-101.
28. Kurre A, van Gool CJ, Bastiaenen CH, Gloor-Juzi T, Straumann D, de Bruin ED. Translation, cross-cultural adaptation and reliability of the german version of the dizziness handicap inventory. *Otol Neurotol.* 2009;30(3):359-67.
29. Gloor-Juzi T, Kurre A, Straumann D, de Bruin ED. Translation and validation of the vertigo symptom scale into German: A cultural adaption to a wider German-speaking population. *BMC Ear Nose Throat Disord.* 2012;12:7.
30. Yagi C, Morita Y, Kitazawa M, Nonomura Y, Yamagishi T, Ohshima S, et al. A Validated Questionnaire to Assess the Severity of Persistent Postural-Perceptual Dizziness (PPPD): The Niigata PPPD Questionnaire (NPQ). *Otol Neurotol.* 2019;40(7):e747-e52.
31. Dunlap PM, Holmberg JM, Whitney SL. Vestibular rehabilitation: advances in peripheral and central vestibular disorders. *Curr Opin Neurol.* 2019;32(1):137-44.
32. Schaaf H, Hesse G. Patients with long-lasting dizziness: a follow-up after neurotological and psychotherapeutic inpatient treatment after a period of at least 1 year. *Eur Arch Otorhinolaryngol.* 2015;272(6):1529-35.
33. Best C, Tschann R, Stieber N, Beutel ME, Eckhardt-Henn A, Dieterich M. STEADFAST: Psychotherapeutic Intervention Improves Postural Strategy of Somatoform Vertigo and Dizziness. *Behav Neurol.* 2015;2015:456850.
34. Schädler S. Schwindel in Supermarkt und Menschenmengen - Fallbeispiel zu den Effekten eines optokinetischen Trainings. *pt Zeitschrift für Physiotherapeuten.* 2020(08):42-5.
35. Schädler S. Der Einfluss eines optokinetischen Trainings auf Schwindel im Supermarkt, in Menschenmengen und durch visuelle Reize – eine Fallbeschreibung *Forum HNO.* 2020(04).
36. Jauregui-Renaud K, Villanueva Padron LA, Cruz Gomez NS. The effect of vestibular rehabilitation supplemented by training of the breathing rhythm or proprioception exercises, in patients with chronic peripheral vestibular disease. *J Vestib Res.* 2007;17(1):63-72.
37. Yardley L, Donovan-Hall M, Smith HE, Walsh BM, Mullee M, Bronstein AM. Effectiveness of primary care-based vestibular rehabilitation for chronic dizziness. *Ann Intern Med.* 2004;141(8):598-605.